

Medications to Avoid Before and After Surgery

If you are taking any medications on this list, they should be discontinued 10-14 days prior to surgery and only Tylenol should be taken for pain. All other medications that you are currently taking must be specifically cleared by your doctor prior to surgery. It is absolutely necessary that all of your current medications be specifically cleared by your doctor and the nursing staff.

Aspirin Medications to Avoid

4-Way Cold Tabs
Adprin-B products
Anacin products
Arthra-G
Arthritis Pain Formula
ASA
Aspergum
Azdone
Backache Maximum Strength Relief
Bismatrol products
Buffetts 11
Butalbital Compound
Cheracol
Cope
Damason-P
Dipentum
Dolobid
Easprin
Equagesic
Fiorinal products
Gensan
Isollyl Improved
Lortab ASA
Magnesium Salicylate
Marthritic
Methocarbamol
Mobigesic
Night-Time Effervescent Cold
Olsalazine
Pabalate products
Panasal
Percodan products
Propoxyphene Compound products
Roxeprin
Salicylate products

5-Aminosalicylic Acid
Alka-Seltzer products
Anexsia w/Codine
Arthriten products
Arthritis Strength BC Powder
Asacol
Asprimox products
Azulfidine products
Bayer Products
Buffered Aspirin
Buffex
Cama Arthritis Pain Reliever
Choline Magnesium Trisalicylate
Coricidin
Darvon Compound-65
Disalcid
Dristan
Ecotrin products
Excedrin products
Gelpirin
Goody's Extra Strength Headache Powders
Kaodene
Magan
Magsal
Meproamate
Micrainin
Momentum
Norgesic products
Orphengesic products
P-A-C
Pentasa
Phenaphen/Codeine #3
Robaxisal
Saleto products
Salsalate

Acetilsalicylic Acid
Amigesic
Argesic-SA
Arthritis Foundation products
Arthropan
Ascriptin products
Axotal
B-A-C
BC Powder
Bufferin products
Butal/ASA/Caff
Carisoprodol Compound
Choline Salicylate
Cortisone Medications
Darvon/ASA
Doan's products
Duragesic
Empirin products
Fiorgen PF
Genprin
Halfprin products
Lanorinal
Magnaprin products
Marnal
Mesalamine
Mobidin
Mono-Gesic
Norwich products
Oxycodone
Pain Reliever Tabs
Pepto-Bismol
Pink Bismuth
Rowasa
Salflex
Salsitab